

Stochastic dynamics of adhesion catch-slip bond cluster

F. LIU¹ and Z.-C. OU-YANG^{1,2}

¹ *Center for Advanced Study, Tsinghua University, Beijing, China*

² *Institute of Theoretical Physics, The Chinese Academy of Sciences, P.O.Box 2735 Beijing 100080, China*

PACS. 87.15.By – Structure and bonding.

PACS. 82.20.Uv – Stochastic theories of rate constants.

Abstract. – We present a general stochastic dynamic model of a cluster of biological complexes with fluctuating dissociation and association rates. The master equation has analytical solutions in two limiting cases: the vanishing force with reflecting boundary condition and slow reaction limit. In particular, the latter can describe the dynamics of a catch-slip bond cluster under a constant force. We find that if the rebinding rate vanishes, the size of the cluster matters little to the cluster lifetime within the whole catch force regime. The large cluster rapidly decays and single catch bond governs the final dynamic behaviors of the cluster. The catch behavior would be further amplified in the presence of large nonvanishing rebinding rates.

Introduction. – Recent experiments found that at single molecule level [1–3], the dissociation rates of some biological adhesion complex, including the bonds between L- and P-selectin and P-selectin glycoprotein ligand 1 (PSGL-1) firstly decrease with initial application of small force, and then increase when the force increases beyond some threshold. This counterintuitive behavior has been termed catch-slip bond transitions. Because these bonds are primarily responsible for the tethering and rolling of leukocytes on inflamed endothelium under shear stress [4, 5], this discovery may provide a direct experimental evidence at the single-molecule level to account for the shear threshold effect [6, 7], in which the number of rolling leukocytes first increases and then decreases while monotonically increasing shear stress. Several kinetic models [2, 8, 9] have been developed to explain the catch-slip bond behavior at the single molecule level. We [10–12] recently suggested alternative mechanism. We proposed that the forced dissociations of PSGL-1–L- and –P might a typical rate process with dynamic disorder [13]. The catch-slip bond transitions can be induced by a particular energy barrier shape with respect to the inner conformational coordinate [11]. Although applied force projection on the dissociation reaction only accelerates the dissociations, the component of the force along the conformational coordinate could stabilize the molecular complex by dragging the molecule to higher energy barriers; if effect of the latter is greater than that of the former, the bond presents catch behavior, otherwise a slip bond is observed.

But before we apply the concept of the catch-slip bond transitions to the real problems, we must know that, in physiological setting adhesion complexes usually exist in a cluster and cooperatively realize their functions with clusters [14]. Two new features will emerge in a

cluster of bonds. Firstly the effect of bond association has to be considered. For single bond case, in general this effect could be neglected due to elastic recoil of the force transducer after complex rupture [15]. In contrast, the dissociated complex can be formed again when the other bonds are still closed. On the other hand, because the closed complexes share the same external force, any dissociation or association of one bond would alter the partition of the force on the other closed bonds. It means that we must consider the cooperativity between bonds in clusters. There are considerable theoretical works contributed to the behaviors of a cluster of slip bonds [14, 16–24]. We naturally ask what difference will be observed between a cluster of catch-slip bonds and slip bonds. We try to give an answer in the present work. Considering that the dissociation of a catch-slip bond is a rate process with dynamic disorder, we begin with a general stochastic dynamical model of a cluster of molecular complex with fluctuating dissociation and association rates.

Model. – Following the previous theoretical work [11], we assume that the forced dissociation rate of single bond has the Bell expression $k_f^+(x) = k^+(x) \exp(\beta f \xi^\ddagger)$, where x is the complex conformational coordinate [25], $\beta = k_B T$, k_B is the Boltzmann's constant, T is absolute temperature, $\xi^\ddagger$ is a projection of the distance from the bound state to the energy barrier onto the external force f , $k^+(x)$ is the intrinsic dissociation rate and has Arrhenius form, i.e., $k^+(x) \propto \exp[\beta \Delta G^\ddagger(x)]$. The rebinding rate $k^-(x)$ is independent of force [14, 17, 24]. Given a cluster of N parallel catch-slip bonds, if the closed bonds sharing the constant force equally, the probability that i bonds are closed and has particular conformational coordinate x_i at time t then satisfies the following one-step master equation

$$\begin{aligned}
\frac{\partial}{\partial t} P_0(x_1^0 \cdots x_N^0, t) &= \left[\sum_i \mathcal{L}^0(x_i) \right] P_0(x_1^0 \cdots x_N^0, t) + \sum_i k_f^+(x_i) P_1(x_1^0 \cdots x_i^1 \cdots x_N^0, t), \\
\frac{\partial}{\partial t} P_1(x_1^0 \cdots x_i^1 \cdots x_N^0, t) &= \left[\mathcal{L}_f^1(x_i) + \sum_{j \neq i} \mathcal{L}^0(x_j) - k_f^+(x_i) - \sum_{j \neq i} k^-(x_j) \right] P_1(x_1^0 \cdots x_i^1 \cdots x_N^0, t) \\
&\quad + \sum_{j \neq i} k_{f/2}^+(x_j) P_2(x_1^0 \cdots x_j^1 \cdots x_i^1 \cdots x_N^0, t), \\
\frac{\partial}{\partial t} P_m(x_1^{\sigma_1} \cdots x_N^{\sigma_N}, t) &= \left[\sum_{b_i} \mathcal{L}_{f/m}^1(x_{b_i}) + \sum_{u_i} \mathcal{L}^0(x_{u_i}) - \sum_{b_i} k_{f/m}^+(x_{b_i}) \right. \\
&\quad \left. - \sum_{u_i} k^-(x_{u_i}) \right] P_m(x_1^{\sigma_1} \cdots x_N^{\sigma_N}, t) \\
&\quad + \sum_{u_i} k_{f/m+1}^+(x_{u_i}) P_{m+1}(x_1^{\sigma_1} \cdots x_{u_i}^1 \cdots x_N^{\sigma_N}, t) \\
&\quad + \sum_{b_i} k^-(x_{b_i}) P_{m-1}(x_1^{\sigma_1} \cdots x_{b_i}^0 \cdots x_N^{\sigma_N}, t), \\
\frac{\partial}{\partial t} P_N(x_1^1 \cdots x_N^1, t) &= \left[\sum_i \mathcal{L}_{f/N}^1(x_i) - \sum_i k_{f/N}^+(x_i) \right] P_N(x_1^1 \cdots x_N^1, t) \\
&\quad + \sum_i k^-(x_i) P_{N-1}(x_1^1 \cdots x_i^0 \cdots x_N^1, t)
\end{aligned} \tag{1}$$

where u_i and b_i respectively denote the ruptured and closed bond number, the superscript $\sigma_i = 1, 0$ respectively represents i -bond ruptured or not, and $\sum_i^N \sigma_i = m$, $2 \leq m \leq N-1$;

the Fokker-Planck operators are defined by

$$\mathcal{L}_f^1(x) = D \frac{\partial}{\partial x} e^{-\beta V_f(x)} \frac{\partial}{\partial x} e^{\beta V_f(x)}, \text{ and } \mathcal{L}^0(x) = D \frac{\partial}{\partial x} e^{-\beta U_0(x)} \frac{\partial}{\partial x} e^{\beta U_0(x)}, \quad (2)$$

where D is the diffusion coefficient, and the potential $V_f(x) = U_1(x) - fx$ [11]. Because rebinding of the completely dissociated state is usually prevented by elastic recoil of the transducer, we write the first two equations ($m=0, 1$) independently. According to previous theoretical work [24], they were termed as the absorbing boundary conditions of the master equation at $m = 0$ and 1. If at initial time, all bonds are closed and in thermal equilibrium, i.e.,

$$P_N(x_1^1, \dots, x_N^1, 0) = \prod_{i=1}^N P_{\text{eq}}^1(x_i), \quad (3)$$

and others vanish. where the distribution $P_{\text{eq}}^1(x_i) \propto \exp[-\beta U_1(x)]$, we'd like to calculate the cluster lifetime, $T_f = \int_0^\infty t D_f(t) dt$. Here

$$D_f(t) = \int dP_0(x_1^0 \dots x_N^0, t) / dt \prod_{i=1}^N dx_i \quad (4)$$

is an extended cluster dissociation rate with dynamic disorder. We see this master equation are far complicated than those in the absence of dynamic disorder. It is impossible to analytically solve eq. (1) for general rate functions and potentials with respect to the conformational coordinate. Numerical approach such as Monte Carlo simulations [24] is also not simple to be applied in the current case. In the following sections, we consider only two limiting cases, the vanishing force with reflecting boundary conditions and slow reaction limits, in which the master equation has analytical solutions.

Vanishing force with reflecting boundary conditions. – In this case, the reflecting boundary condition [24] means that the first two equations in eq. (1) are altered into,

$$\begin{aligned} \frac{\partial}{\partial t} P_0(x_1^0 \dots x_N^0, t) &= \left[\sum_i \mathcal{L}^0(x_i) - \sum_i k^-(x_i) \right] P_0(x_1^0 \dots x_N^0, t) \\ &\quad + \sum_i k^+(x_i) P_1(x_1^0 \dots x_i^1 \dots x_N^0, t), \\ \frac{\partial}{\partial t} P_1(x_1^0 \dots x_i^1 \dots x_N^0, t) &= \left[\mathcal{L}^1(x_i) + \sum_{j \neq i} \mathcal{L}^0(x_j) - k^+(x_i) \right. \\ &\quad \left. - \sum_{j \neq i} k^-(x_j) \right] P_1(x_1^0 \dots x_i^1 \dots x_N^0, t) \\ &\quad + \sum_{j \neq i} k^+(x_j) P_2(x_1^0 \dots x_j^1 \dots x_i^1 \dots x_N^0, t) \\ &\quad + k^-(x_i) P_0(x_1^0 \dots x_i^0 \dots x_N^0, t). \end{aligned}$$

The other terms ($m \geq 2$) are the same with those in eq. (1) except the force therein vanishing. It is easy to prove that $P_m(x_1^{\sigma_1} \dots x_N^{\sigma_N}, t) = \prod_{i=1}^N P^{\sigma_i}(x_i, t)$ is the solution in this limit, where

$P^\sigma(x, t)$, $\sigma = 0, 1$ are solutions of the dissociation-association diffusion equations of single bond in the absence of force given by

$$\begin{aligned}\frac{\partial P^1(x, t)}{\partial t} &= \mathcal{L}^1(x)P^1(x, t) - k^+(x)P^1(x, t) + k^-(x)P^0(x, t), \\ \frac{\partial P^0(x, t)}{\partial t} &= \mathcal{L}^0(x)P^0(x, t) + k^+(x)P^1(x, t) - k^-(x)P^0(x, t).\end{aligned}\quad (5)$$

Here the initial conditions are $P^1(x, 0) \propto \exp(-\beta U_1(x))$ and $P^0(x, 0) = 0$. Therefore, any quantity of interest can now be calculated. For instance, the probability $Q_m(t)$ that m -bond are closed at time t is

$$\begin{aligned}Q_m(t) &= \sum_{\sigma_1 \dots \sigma_N} \int P_m(x_1^{\sigma_1} \dots x_N^{\sigma_N}, t) \prod_1^N dx_i \\ &= C_N^m \left(\int P^1(x, t) dx \right)^m \left(\int P^0(x, t) dx \right)^{N-m}.\end{aligned}\quad (6)$$

Although the master equation with reflecting boundary conditions has been formally solved, it is still a great challenge to solve eq. (5) analytically. Various approximation approaches have been developed in the past two decades [26]. We are not ready to further discuss it. Instead, we study the other limit, in which is relevant to the dynamics of a catch-slip bond cluster.

Slow reaction limit. – In this limit the processes of the rupture and rebinding of any single bonds are very slow compared to the conformational diffusions. The thermal equilibrium distribution of the conformation for each bond hence is always maintained during the courses of “reactions”. We assume

$$P_m(x_1^{\sigma_1} \dots x_N^{\sigma_N}, t) = q_m(\sigma_1 \dots \sigma_N, t) \prod_{b_i} P_{\text{eq}}^1(x_{b_i}, f/m) \prod_{u_i} P_{\text{eq}}^0(x_{u_i}), \quad (7)$$

where P_{eq}^σ , $\sigma = 0, 1$ are the thermal equilibrium distributions and satisfy

$$\mathcal{L}_{f/m}^1(x_{b_i})P_{\text{eq}}^1(x_{b_i}, f/m) = 0, \text{ and } \mathcal{L}^0(x_{u_i})P_{\text{eq}}^0(x_{u_i}) = 0, \quad (8)$$

respectively. Substituting in eq. (1) and integrating all conformational freedoms lead to a master equation about $q_m(\sigma_1 \dots \sigma_N, t)$. According to the definition of $Q_m(t)$ (eq. (6)), we can further simplify the new master equation as follows,

$$\begin{aligned}\frac{d}{dt}Q_0(t) &= K_1^+(f)Q_1(t) \\ \frac{d}{dt}Q_1(t) &= -[K_1^+(f) + K_1^-]Q_1(t) + K_2^+Q_2(t) \\ \frac{d}{dt}Q_m(t) &= -[K_m^+(f) + K_m^-]Q_m(t) - K_{m+1}^+(f)Q_{m+1}(t) + K_{m-1}^-Q_{m-1}(t) \\ \frac{d}{dt}Q_N(t) &= -K_N^+(f)Q_N(t) + K_{N-1}^-Q_{N-1}(t)\end{aligned}\quad (9)$$

where

$$\begin{aligned}K_m^+(f) &= m e^{\beta f \xi^\dagger / m} \int k^+(x) P_{\text{eq}}^1(x, f/m) dx, \\ K_m^- &= (N - m) \int k^-(x) P_{\text{eq}}^0(x) dx = (N - m) \langle k^- \rangle.\end{aligned}\quad (10)$$

We see eq. (9) is formally the same with the one-step master equation that describes the force ruptured dynamics of a cluster of slip bond [24]. Many analytical results that are independent of concrete expression of the rates hence could be applied here. On the other hand, because the new defined rate constant $K_m^+(f)$ is a dynamical average of the Bell expression on the distribution $P_{\text{eq}}(x, f/m)$, its force dependence could be very different from that of conventional slip bond. Particularly, if the potential $U_1(x)$ is harmonic with spring constant κ , and barrier $\Delta G^\ddagger(x)$ is a piecewise function with two segments, $K_m^+(f)$ is approximated to be

$$K_m^+(f) \approx \begin{cases} mk_c \exp(-\beta d_c^\ddagger f/m) & f < f_t \\ mk_s \exp(+\beta d_s^\ddagger f/m) & f > f_t, \end{cases}$$

where f_t is the bond transition force, k_i , $i = d, s$ are some constants, and the “effective” distance $d_i^\ddagger > 0$, are the simple combinations of κ , the slopes of segments, and $\xi^\ddagger$ [11]. We see if force is smaller than f_t , the rupture rate $K_m^+(f)$ decreases with increasing force (a catch bond cluster); otherwise it is a slip bond cluster. For the latter, eq. (9) is completely the same with previous models [22, 24]. Hence we only study the dynamics of the catch bond cluster. We respectively discuss two cases according to zero and nonzero rebinding rates.

Rebinding rates $k^-(x) = 0$. In this case, the probability $Q_m(t)$ of m -bond closed at time t has an exact solution given by [22]

$$Q_m(t) = \left(\prod_{n=m+1}^N K_n^+(f) \right) \sum_{n=m}^N \left[e^{-tK_n^+(f)} \prod_{\substack{p=m \\ p \neq n}}^N (K_p^+(f) - K_n^+(f))^{-1} \right] \quad (11)$$

Hence, any quantity of interest can be calculated from $Q_m(t)$. For instance, the cluster lifetime T_f is

$$T_f = \sum_{m=1}^N K_m^+(f)^{-1}. \quad (12)$$

For large N and small force $f < f_c = k_B T / d_c$, the above equation can be approximated to be [28]

$$T_f \approx H_N \approx \ln N + 1/2N + 0.577, \quad (13)$$

where $H_N = \sum_{i=1}^N i^{-1}$ is the harmonic number. It is same with the lifetime of slip bond cluster [24]. It is expected because at small forces the difference between the catch and slip bonds are negligible. If $f > f_c$, the force in the exponentials of T_f cannot be neglected again. In contrast, for the larger force the dominating term in eq. (12) is the one for the rupture of only one bond, i.e.,

$$T_f \sim k_c^{-1} \exp(f/f_c). \quad (14)$$

This result is further demonstrated by the time decay function of the mean number of the closed bonds on large time scales:

$$\langle m \rangle \sim \exp[-tk_c \exp(-f/f_c)]. \quad (15)$$

Compared to the behaviors of a slip bond cluster with large size, three difference are observed in a large size of catch bond cluster if the rebinding rate vanishes. First, the size of the

catch bond cluster matters little for the cluster lifetime within the whole catch force regime, whereas it only holds for smaller forces in the slip bond cluster. Then, for larger force the dynamics of the catch cluster is took over by single catch bond. In contrast, the largest clusters dominate the decay of the slip bond cluster [24]. Finally, as a characteristic of a catch bond, the increasing force only exponentially elongates the lifetime of the catch bond cluster.

Nonzero rebinding rates. Although $Q_m(t)$ in Eq. (9) in principle could be calculated by eigenvalue technique in this case, except the simplest case $N = 2$, it is not more efficient than the exact stochastic simulations [24]. Fortunately, it has been proved [24, 27] that the cluster lifetime of great interest has still an exact form

$$T_f = \sum_{m=1}^N K_m^+(f)^{-1} + \sum_{m=1}^{N-1} \sum_{n=m+1}^N \left(\prod_{p=n-m}^{n-1} K_p^- / \prod_{p=n-m}^n K_p^+(f) \right). \quad (16)$$

We are not interested in the general dependence of eq. (16) on force and N . Instead we consider a specific case where the ratio $\gamma = \langle k^- \rangle / k_c$ is larger than 1. Thus for large N and force, eq. (16) is

$$T_f \sim k_c^{-1} e^{f/f_c} \left[1 + \frac{1}{N} \gamma^{N-1} e^{f(H_N-1)/f_c} \right]. \quad (17)$$

In contrast to the case without rebinding, the size of the catch cluster will change the cluster lifetime dramatically as $\gamma > 1$; the presence of the rebinding rate further amplifies the lifetime by γ^{N-1} -fold. Therefore, the effect of the rebinding are the same for the slip and catch bond cluster.

Discussion and conclusions. – In this work, we present a general stochastic dynamic model of a cluster of biological complexes with dynamic disorder. We study two simplest limiting cases: vanishing force with reflecting boundary conditions and the slow reaction limits. Under the former limit, because the dissociation and association of each bonds are independent, the solution to the model is a simple linear combination of the dynamics of single bond. Under the slow reaction limit, we demonstrate that the model can be reduced to the conventional one-step master equation except that the rates are replaced by dynamic averages. Particularly, given a specific energy barrier with respect to molecular conformational coordinate, this master equation can describe the dynamics of a cluster of catch-slip bonds. We found that many conclusions obtained for the slip bond cluster could be applied to the catch bond cluster with minor modification. We give simple relations between the cluster lifetime and the cluster size N , force f and the average rebinding rate. These conclusion would be useful in future dynamic studies of the catch-slip bond clusters.

* * *

One of the authors (F.L.) would like to thank Prof. Seifert, U. for providing his interesting works to us. F.L. was supported by the National Natural Science Foundation of China (NSFC).

REFERENCES

- [1] MARSHALL, B. T., LONG, M., PIPER, J. W., YAGO, T., MCIVER, R. P. and ZHU, C., *Nature*, **423** (2003) 190.
- [2] EVANS, E., LEUNG, A., HEINRICH, V. and ZHU, C., *Proc. Natl. Acad. Sci. USA*, **101** (11281) 2004.

- [3] SARANGAPANI, K. K., YAGO, T., KLOPOCKI, A. G., LAWRENCE, M. B., FIEGER, C. B., ROSEN, S. D., MCEVER, R. P. and ZHU, C., *J. Biol. Chem.*, **279** (2003) 2291.
- [4] MCEVER, R. P., *Curr. Opin. Cell Biol.*, **14** (2002) 581.
- [5] KOONSTANTOPOULOS, K., KURKRETI, S. and MCINTIRE, L. V., *Adv. Drug Deliv. Rev.*, **33** (1998) 141.
- [6] FINGER, E. B., PURI, K. D., ALON, R., LAWRENCE, M. B., VON ANDRIAN, U. H. and SPRINGER, T. A., *Nature*, **279** (1996) 266.
- [7] LAWRENCE, M. B., KANSAS, G. S., KUNKEL, E. J. and LEY, K., *J. Cell. Biol.*, **136** (1997) 717.
- [8] BARSEGOV, V. and THIRUMALAI, D., *Proc. Natl. Acad. Sci. USA*, **102** (2005) 1835.
- [9] PEREVERZEV, Y. V., PREZHDO, O. V., FORERO, M., SOKURENKO, E. V. and THOMAS, W., *Biophys. J.*, **89** (2005) 1446.
- [10] LIU, F., OU-YANG, Z.-C. and IWAMOTO, M., *Phys. Rev. E*, **73** (2006) 010901(R).
- [11] LIU, F. and OU-YANG, Z.-C., *Force Modulating Dynamic Disorder: A Physical Model of Catch-slip Bond Transitions in Receptor-Ligand Forced Dissociation Experiments*, submitted to *Phys. Rev. E*, q-bio.CB/0601017 preprint, 2006.
- [12] LIU, F. and OU-YANG, Z.-C., *Bell Rate Model with Dynamic Disorder: Model and Its Application in the Receptor-ligand Forced Dissociation Experiments*, cond-mat/0604499 preprint, 2006.
- [13] ZWANZIG, R., *Acc. Chem. Res.*, **23** (1990) 148.
- [14] BELL, G. I., *Science*, **200** (1978) 618.
- [15] EVANS, E., *Annu. Rev. Biophys. Biomol. Struct.*, **30** (2001) 105.
- [16] DEMBO, M., TOURNEY, D. C., SAXMAN, K. and HAMMER, D., *Proc. R. Soc. Lond. B*, **234** (1988) 55.
- [17] SEIFERT, U., *Phys. Rev. Lett.*, **84** (2000) 2750.
- [18] SEIFERT, U., *Europhys. Lett.*, **58** (2002) 792.
- [19] LIPOWSKY, R., *Phys. Rev. Lett.*, **77** (1996) 1652.
- [20] TEES, D. F. J., WOODWARD, J. T., and HAMMER, D. A., *J. Chem. Phys.*, **114** (2001) 7483.
- [21] COZENS-ROBERTS, C., LAUFFENBURGER, D. A. and QUINN, J. A., *Biophys. J.*, **58** (1990) 841.
- [22] ERDMANN, T. and SCHWARZ, U. S., *Phys. Rev. Lett.*, **92** (2004) 108102.
- [23] ERDMANN, T. and SCHWARZ, U. S., *Europhys. Lett.*, **66** (2004) 603.
- [24] ERDMANN, T. and SCHWARZ, U. S., *J. Chem. Phys.*, **121** (2004) 8997.
- [25] AGMON, N. and HOPFIELD, J. J., *J. Chem. Phys.*, **78** (1983) 6947.
- [26] PORTMAN, J. J. and WOLYNES, P. G., and references therein, *J. Phys. Chem. A*, **103** (1999) 10602.
- [27] VAN KAMPEN, N. G., *Stochastic Processes in Physics and Chemistry* (Elsevier, Amsterdam) 1992.
- [28] GOLDSTEIN B. and WOFSY, C., *Immunol. Today*, **17** (1996) 77.